

Thermistor data at 10 cm intervals in air, ice, water from autonomous buoys from ARSV Laurence M. Gould LMG0106 in the Southern Ocean, August to November, 2001 (SOGLOBEC project)

Website: <https://www.bco-dmo.org/dataset/3115>

Data Type: Cruise Results

Version: 1

Version Date: 2009-05-07

Project

» [U.S. GLOBEC Southern Ocean](#) (SOGLOBEC)

Program

» [U.S. GLOBal ocean ECosystems dynamics](#) (U.S. GLOBEC)

Contributors	Affiliation	Role
Costa, Daniel P.	University of California-Santa Cruz (UC Santa Cruz)	Principal Investigator
Fritsen, Chris H.	Desert Research Institute (DRI)	Co-Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Thermistor data at 10 cm intervals in air, ice, water from autonomous buoys from ARSV Laurence M. Gould LMG0106 in the Southern Ocean, August to November, 2001 for the Southern Ocean GLOBEC project.

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
- [Parameters](#)
- [Instruments](#)

- [Deployments](#)
 - [Project Information](#)
 - [Program Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: N:-67 E:-68 S:-69.5 W:-75

Temporal Extent: 2001-08-06 - 2001-10-11

Dataset Description

Related datasets:

[buoy_position](#) Autonomous buoy position data, Aug. - Nov. 2001

[buoy_temp_barom](#) Air temperature and barometric pressure results from autonomous buoys, Aug. - Nov. 2001

[buoy_radm](#) Radiometer data from autonomous buoy 07949, Aug. - Nov. 2001

Acquisition Description


During the July-September 2001 Southern Globec cruises of the Palmer and Gould we installed four buoys. Two of the buoys (07413, 07440) were standard off-the-shelf Met-Ocean Ice buoys (see above photograph). These buoys measured barometric pressure, air temperature, and GPS position. The data were transmitted via ARGOS. A radar reflector was mounted on a wooden 4 x 4 to help find the site if there was an opportunity to return, however the reflector return was difficult to discern among all the backscatter clutter caused by deformed ice and icebergs.

The other two buoys (07949, 07950) were custom made CRREL ice mass balance buoys (photo on right). These buoys reported barometric pressure, Argos position, and air temperature. In addition they had a thermistor string that measured a vertical profile of temperature, at 10 cm spacing, from the air, through the snow and ice, and into the upper ocean. There were acoustic sensors measuring the positions of the snow surface and ice bottom. A fluorometer was mounted under the ice. One of the buoys (07949) had three spectroradiometers; one mounted about the ice, one mounted directly below the ice, and one a few meters deep in the upper ocean. Results from all buoys are compared.

Positive values are above the ice; negative values are in the ice and underlying ocean. Due to surface flooding and freezing, the position of the snow-ice interface changed during the

measurement period.

Tracks from all 2001 buoys:



[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
therm_pos	height of thermistor relative to ice. Positive values are above the ice; negative values are in the ice and underlying ocean.	centimeters
yrday_gmt	GMT day and decimal time, as 326.5 for the 326th day of the year, or November 22 at 1200 hours (noon)	
date_gmt	gmt month, day and year, usually as a text string, e.g. feb10_1995.	
year	year, reported as YYYY, e.g. 1995	
month_gmt	month of year, GMT time , i.e. 01-12	
day_gmt	day, GMT time e.g. 22.	
time_gmt	time of day, reported in GMT time, 24 hour clock	
temp	temperature at measurement depth	degrees Celsius

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Drifter Buoy
Generic Instrument Name	Drifter Buoy
Dataset-specific Description	4 buoys in ice
Generic Instrument Description	<p>Drifter buoy to include the Beardsley Drifter. Generic drifter buoys may be surface or sub-surface buoys that move with the current. They have a variety of instruments attached, providing a platform that allows for the measurement of surface drifts, air pressure and other variables. The Beardsley Drifters are near-surface satellite-tracked drifters used for observations of circulation patterns. They are WOCE-style drifters featuring holey sock drogues. Each drifter has a small (~ 30 cm diameter) surface float with ARGOS transmitter and batteries tethered to a holey sock drogue centered at 15 m below the surface. The drogue, about 10 m tall and 1 m in diameter, is designed to "lock" itself to the water so that the surface float follows the mean water motion at 15 m depth with very little slippage even in high winds. Thus measuring the drifter's position as a function of time provides a Lagrangian measurement of the 15-m ocean current.</p> <p>(http://globec.who.edu/jg/info/globec/soglobec/drifters_argos%7Bdir=glob...?)</p> <p>WOCE-drifters: http://woce.nodc.noaa.gov/wdiu/diu_summaries/svp/index.htm</p>

[[table of contents](#) | [back to top](#)]

Deployments

LMG0106

Website	https://www.bco-dmo.org/deployment/57639
Platform	ARSV Laurence M. Gould
Report	http://www.ccpo.odu.edu/Research/globec/cruises01/lmg0106_menu.html
Start Date	2001-07-21
End Date	2001-09-01

Project Information

U.S. GLOBEC Southern Ocean (SOGLOBEC)

Website: http://www.ccpo.odu.edu/Research/globec_menu.html

Coverage: Southern Ocean

The fundamental objectives of United States Global Ocean Ecosystems Dynamics (U.S. GLOBEC) Program are dependent upon the cooperation of scientists from several disciplines. Physicists, biologists, and chemists must make use of data collected during U.S. GLOBEC field programs to further our understanding of the interplay of physics, biology, and chemistry. Our objectives require quantitative analysis of interdisciplinary data sets and, therefore, data must be exchanged between researchers. To extract the full scientific value, data must be made available to the scientific community on a timely basis.

Program Information

U.S. GLOBal ocean ECosystems dynamics (U.S. GLOBEC)

Website: <http://www.usglobec.org/>

Coverage: Global

U.S. GLOBEC (GLOBal ocean ECosystems dynamics) is a research program organized by oceanographers and fisheries scientists to address the question of how global climate change may affect the abundance and production of animals in the sea. The U.S. GLOBEC Program currently had major research efforts underway in the Georges Bank / Northwest Atlantic Region, and the Northeast Pacific (with components in the California Current and in the Coastal Gulf of Alaska). U.S. GLOBEC was a major contributor to International GLOBEC efforts in the Southern Ocean and Western Antarctic Peninsula (WAP).

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Antarctic Sciences (NSF ANT)	unknown SOGLOBEC NSF ANT

[[table of contents](#) | [back to top](#)]